

nautics and stration

ce Flight Center ryland 20771

#### The NASA STI Program Office ... in Profile

Since its founding, NASA has been dedicated to the advancement of aeronautics and space science. The NASA Scientific and Technical Information (STI) Program Office plays a key part in helping NASA maintain this important role.

The NASA STI Program Office is operated by Langley Research Center, the lead center for NASA's scientific and technical information. The NASA STI Program Office provides access to the NASA STI Database, the largest collection of aeronautical and space science STI in the world. The Program Office is also NASA's institutional mechanism for disseminating the results of its research and development activities. These results are published by NASA in the NASA STI Report Series, which includes the following report types:

- TECHNICAL PUBLICATION. Reports of completed research or a major significant phase of research that present the results of NASA programs and include extensive data or theoretical analysis. Includes compilations of significant scientific and technical data and information deemed to be of continuing reference value. NASA's counterpart of peer-reviewed formal professional papers but has less stringent limitations on manuscript length and extent of graphic presentations.
- TECHNICAL MEMORANDUM. Scientific and technical findings that are preliminary or of specialized interest, e.g., quick release reports, working papers, and bibliographies that contain minimal annotation. Does not contain extensive analysis.
- CONTRACTOR REPORT. Scientific and technical findings by NASA-sponsored contractors and grantees.

- CONFERENCE PUBLICATION. Collected papers from scientific and technical conferences, symposia, seminars, or other meetings sponsored or cosponsored by NASA.
- SPECIAL PUBLICATION. Scientific, technical, or historical information from NASA programs, projects, and mission, often concerned with subjects having substantial public interest.
- TECHNICAL TRANSLATION.
   English-language translations of foreign scientific and technical material pertinent to NASA's mission.

Specialized services that complement the STI Program Office's diverse offerings include creating custom thesauri, building customized databases, organizing and publishing research results . . . even providing videos.

For more information about the NASA STI Program Office, see the following:

- Access the NASA STI Program Home Page at http://www.sti.nasa.gov/STI-homepage.html
- E-mail your question via the Internet to help@sti.nasa.gov
- Fax your question to the NASA Access Help Desk at (301) 621-0134
- Telephone the NASA Access Help Desk at (301) 621-0390
- Write to: NASA Access Help Desk

NASA Access Help Desk NASA Center for AeroSpace Information 7121 Standard Drive Hanover, MD 21076-1320

## NASA/TM—2000–209891, Vol. 219



# Technical Report Series on the Boreal Ecosystem-Atmosphere Study (BOREAS)

Forrest G. Hall and Sara K. Conrad, Editors

# Volume 219 BOREAS TGB-1 NSA CH<sub>4</sub> and CO<sub>2</sub> Chamber Flux Data

Patrick Crill and Ruth K. Varner University of New Hampshire, Durham

National Aeronautics and Space Administration

Goddard Space Flight Center Greenbelt, Maryland 20771

	A!1-1-1 C	
	Available from:	
NASA Center for AeroSpace Information		National Technical Information Service
7101 G. 1 1D.		
7121 Standard Drive		5285 Port Royal Road
Hanover, MD 21076-1320		Springfield, VA 22161
Price Code: A17		Price Code: A10
THE COUE, AT		rnce Coue. A10

## BOREAS TGB-1 NSA CH4 and CO2 Chamber Flux Data

Patrick M. Crill, Ruth K. Varner

## Summary

The BOREAS TGB-1 team made methane (CH<sub>4</sub>) and carbon dioxide (CO<sub>2</sub>) dark chamber flux measurements at the NSA-OJP, NSA-OBS, NSA-BP, and NSA-YJP sites from 16-May-1994 through 13-Sep-1994. Gas samples were extracted approximately every 7 days from dark chambers and analyzed at the NSA lab facility. The data are provided in tabular ASCII files.

#### **Table of Contents**

- 1) Data Set Overview
- 2) Investigator(s)
- 3) Theory of Measurements
- 4) Equipment
- 5) Data Acquisition Methods
- 6) Observations
- 7) Data Description
- 8) Data Organization
- 9) Data Manipulations
- 10) Errors
- 11) Notes
- 12) Application of the Data Set
- 13) Future Modifications and Plans
- 14) Software
- 15) Data Access
- 16) Output Products and Availability
- 17) References
- 18) Glossary of Terms
- 19) List of Acronyms
- 20) Document Information

#### 1. Data Set Overview

#### 1.1 Data Set Identification

BOREAS TGB-01 NSA CH<sub>4</sub> and CO<sub>2</sub> Chamber Flux Data

#### 1.2 Data Set Introduction

Chamber flux measurements were taken at the Northern Study Area (NSA) Old Jack Pine (OJP), Young Jack Pine (YJP), Old Black Spruce (OBS), and Beaver Pond (BP) sites during the summer of 1994 by the BOReal Ecosystem-Atmosphere Study (BOREAS) Trace Gas Biogeochemistry (TGB)-01 team. The values represent the average/mean flux from five values taken at 20-minute intervals.

#### 1.3 Objective/Purpose

The purpose of these measurements was to examine the trace gas exchange between the atmosphere and the boreal soils.

#### 1.4 Summary of Parameters

CH<sub>4</sub> and CO<sub>2</sub> chamber fluxes were measured at the NSA OJP, YJP, OBS, and BP sites. Temperatures of the chamber, 1-cm soil depth, and 10-cm soil depth were recorded at the OJP, YJP, and OBS (Patrick Crill) sites. At the OBS (Dean Moosavi) and BP sites, temperatures at 5-, 10-, and 20-cm soil depths were recorded.

#### 1.5 Discussion

The chamber flux data are provided in American Standard Code for Information Interchange (ASCII) comma-delimited text files. The data are a compilation of measurements from the aluminum chamber and collar sites at OJP, YJP, and OBS (Patrick Crill) and the plastic chamber and collar sites at OBS (Dean Moosavi) and BP sites. Another part of the data is a diurnal study completed 08-Jul-1994 and 09-Jul-1994 at the BP plastic collar and chamber sites.

#### 1.6 Related Data Sets

BOREAS TGB-01 CH4 Tower flux data over NSA BOREAS TGB-01 Soil CH4 and CO2 concentration data over the NSA

## 2. Investigator(s)

#### 2.1 Investigator(s) Name and Title

Dr. Patrick M. Crill Research Associate Professor University of New Hampshire

#### 2.2 Title of Investigation

Magnitude and Control of Trace Gas Exchange in Boreal Ecosystems

#### 2.3 Contact Information

#### Contact 1:

Dr. Patrick M. Crill Institute for the Study of Earth, Oceans, and Space Complex Systems Research Center University of New Hampshire Durham, NH 03824 (603) 862-3519 (603) 862-0188 (fax)

#### Contact 2:

Sadredin C. Moosavi Graduate Student Institute for the Study of Earth, Oceans, and Space Complex Systems Research Center University of New Hampshire Durham, NH 03824 (603) 862-2927 (603) 862-0188 (fax)

#### Contact 3:

Ruth K. Varner Research Scientist Institute for the Study of Earth, Oceans, and Space Complex Systems Research Center University of New Hampshire Durham, NH 03824 (603) 862-2939 (603) 862-0188 (fax)

#### Contact 4:

Jeffrey A. Newcomer Raytheon ITSS Code 923 NASA GSFC Greenbelt, MD 20771 (301) 286-7858 (301) 286-0239 (fax) Jeffrey.Newcomer@gsfc.nasa.gov

## 3. Theory of Measurements

Chamber fluxes measure the changes in the mixing ratio of trace gases (CO<sub>2</sub> and CH<sub>4</sub>) in a closed headspace over a period of time. This headspace is isolated from the atmosphere; therefore, the exchange of material between the covered soil and the headspace can be quantified.

## 4. Equipment

#### 4.1 Sensor/Instrument Description

CH<sub>4</sub> was quantified with a Shimadzu GC-14A Gas Chromatograph (GC) or a Shimadzu GC-MINI2 with a flame ionization detector (FID) operated at 125 °C after separation on a HayeSepQ column at 40 °C using ultrapure (99.999%) N<sub>2</sub> as a carrier gas flowing at 30 mL/min. CO<sub>2</sub> was quantified with a Shimadzu GC-14A with a thermal conductivity detector (TCD) operated at 70 °C after separation on a HayeSepQ column at 40 °C using ultrapure (99.999%) He as a carrier gas flowing at 30 mL/min. The oven temperature during sample analysis was run at 130 °C and the detector at 300 °C. Analog signals (0-1 V) from the detectors were digitized at 10 Hz with a Hewlett Packard (HP) 35000D A/D board and quantified and logged using HP ChemStation software. Chamber fluxes were accomplished with aluminum chambers manufactured at the University of New Hampshire and designed by Patrick Crill.

#### 4.1.1 Collection Environment

The chamber fluxes were collected under ambient conditions. The GC analysis was completed at the Hayes Road Lab in Thompson, Manitoba.

#### 4.1.2 Source/Platform

Ground.

#### 4.1.3 Source/Platform Mission Objectives

The mission objective was to determine the soil-surface exchange rates of CH<sub>4</sub> and CO<sub>2</sub> at the NSA OJP, YJP, OBS, and BP sites.

#### 4.1.4 Key Variables

The key variables measured during the sampling period were CH<sub>4</sub> and CO<sub>2</sub> fluxes. The OBS site was divided into two different sets of data. In the SITE\_COLLAR, a C denotes the OBS site where Patrick Crill measured fluxes with aluminum chambers and collars. An M denotes the OBS site where Dean Moosavi measured fluxes with plastic chambers and collars. Plastic chambers and collars were also used at the BP site. Soil profile temperatures were also recorded.

#### 4.1.5 Principles of Operation

The Shimadzu GC-14A GC is equipped with a hydrogen FID and a TCD. The FID is used to detect CH<sub>4</sub>, while the TCD is used to detect CO<sub>2</sub>. The FID uses a hydrogen flame in an air atmosphere to burn components as they exit the column. In the flame, carbon-carbon bonds are fragmented so that various organic ions and free electrons exist. Application of a voltage across a collector electrode over the flame causes an ion current to flow, which is amplified and then measured as the output signal.

The TCD detects CO<sub>2</sub> by passing a sample, in a helium carrier gas, past metallic filaments with current flowing through them. The sample components with lower thermal conductivity than the helium carrier gas raise the filament temperature when they pass through. The signal output from the TCD is a measurement of the change in filament resistance caused by the temperature rise. The signal output from both the FID and TCD is for a data processor, integrator, recorder, or computer (Instruction Manual: GC-14A; Shimadzu Corporation, Kyoto, Japan). The GC-MINI2 was equipped with a FID and operated in the same manner as the GC-14A FID.

#### 4.1.6 Sensor/Instrument Measurement Geometry

Not applicable.

#### 4.1.7 Manufacturer of Sensor/Instrument

Manufacturer of GC-14A FID/TCD, GC-MINI2, and GC-8A ECD Shimadzu Scientific Instruments, Inc. 7102 Riverwood Drive Columbia, MD 21046 (410) 381-1227

#### 4.2 Calibration

#### 4.2.1 Specifications

None given.

#### 4.2.1.1 Tolerance

The sensitivity of the TCD is approximately 6000 mV ml/mg. The FID's maximum sensitivity is 3 x  $10^{-12}$  g/s for diphenyl.

#### 4.2.2 Frequency of Calibration

The instrument is calibrated on a daily basis. Standards are run generally before and after samples on a given day of analysis.

#### 4.2.3 Other Calibration Information

Signal peaks from the detectors were quantified with working standards calibrated against Canadian Atmospheric Environment Services (AES) certified primary standards acquired by the BOREAS project and a CO<sub>2</sub>/CH<sub>4</sub> standard of Niwot Ridge air prepared by the National Oceanic and Atmospheric Administration (NOAA) Climate Monitoring and Diagnostics Laboratory (CMDL). Uncertainty in the standards' analyses on a given day ranged from 0.1 to 0.2%.

## 5. Data Acquisition Methods

The chamber fluxes are determined by analysis of concentrations of methane (CH<sub>4</sub>) and carbon dioxide (CO<sub>2</sub>) in a time series of grab samples of headspace over the ground surface enclosed by a dark aluminum chamber. The volumes of the two aluminum chambers were 0.071 and 0.120 m³ over an area of 0.397 m². The aluminum chamber was placed on the trough of an aluminum collar embedded in the ground. Water was added to the trough of the collar to create an airtight seal. The volume of the plastic chambers was 0.028 m³ over an area of 0.078 m². The seal between chamber and collar was made using weather stripping or water as appropriate. Five 60-mL samples were removed from the headspace with polypropylene syringes and polycarbonate/nylon stopcocks at 4-minute intervals for 20 minutes (five samples). Samples were returned to the Hayes Road lab and analyzed for CO<sub>2</sub> and CH<sub>4</sub> using gas chromatography within 12 hours after collection.

#### 6. Observations

#### 6.1 Data Notes

None given.

#### **6.2 Field Notes**

#### For data collected by Patrick Crill:

- 25-May-94- Fans in chambers not working at OBS. Can't use the data.
- 29-May-94- Fan in chamber not working after 20 minutes in OJPA2 flux; no fan for the OJPM2 flux; no fan for the OJP1 and 2 fluxes.
- 01-Jun-94- No fan for OBS12 flux.
- 20-Jun-94- Thermometer probe outside of chamber for collar OBS1.
- 10-Jul-94- One chamber had been filled with water and therefore the fan and temperature probe aren't working. Used it with collars YJP1, OJPA1, OJPL2, OJPM1, and OJPM2.
- 16-Jul-94- No temperature still in one chamber at OJP and YJP.
- 23-Jul-94- No temperature still in one chamber at OJP and YJP; Collar YJP2 syringe number 4 lost its stopcock.
- 30-Jul-94- No temperature still in one chamber at OJP and YJP.
- 01-Aug-94- Very hot, using silver mylar to reflect sun on half of the flux chambers to see if it effects temperature change in chamber...it doesn't heat up as much as chambers without cover by about 10 °C.
- 07-Aug-94- Fixed chamber thermometer for first few fluxes; then it went bad again.
- 08-Aug-94- No fan for collar OBS10 flux; no temperatures for OBS8, OBS6, OBS11, OBS4, and OBS2
- 12-Aug-94- No temperature still in one chamber at OJP and YJP.
- 19-Aug-94- No temperature still in one chamber at OJP and YJP.

#### For data taken by Dean Moosavi:

- 19-May-94- Experiments halted due to chamber hardware problems.
- 20-May-94- Screw missing in chamber for flux at collar OBS1; therefore, there was a hole in chamber during the flux. Fixed with duct tape.
- 26-May-94- Problems with collar at collar BP29.
- 04-Jun-94- Water levels in the fen have dropped 5-10 cm from season beginning at OBS.
- 10-Jun-94- Carex and other vegetation have sprouted. Larch are out.
- 20-Jun-94- Temperature probe dead!
- 21-Jun-94- Fans did not work for collars 19 and 20 at BP.
- 30-Jun-94- BP- Bear attack on equipment! Collar 23 pulled up and moved 5 yards. Minor tooth damage. Collar 18 destroyed. Lichen thermocouple uprooted and destroyed. Pond and dam level have dropped 10 cm leaving collars above the water level.

- 02-Aug-94- BP- Moose stepped in collar 24!
- 21-Aug-94- Senescence beginning.
- 24-Aug-94- Due to problems with yesterday's samples. BP and dam were resampled off collar in vicinity of site.
- 31-Aug-94- BP and dam fluxes taken next to (NOT on) collars.
- 13-Sep-94- BP collars 25-32 fluxes taken near (not on) collars.
- 14-Sep-94- All collars and chambers removed!

## 7. Data Description

#### 7.1 Spatial Characteristics

#### 7.1.1 Spatial Coverage

Descriptions of the sites and their North American Datum of 1983 (NAD83) coordinates are:

- The collars at the OJP tower site and profile sampling sites were located on moss plots that were approximately northwest of the tower, while the profile corral was located a few meters south of the collars (55°55'43.2" N, 98°37'29.4" W).
- The collars at the OJP site located on lichen plots were due west of the tower; the profile trench was located about 1 m southeast of the collars (55°55'40.9" N, 98°37'29.3" W).
- The collars at the YJP tower site were located southeast of the tower; the gas profile corral was located approximately 1 m west of the collars (55°53'42.8" N, 98°17'12.7" W).
- The collars at the OBS tower site were located along a low to high moisture gradient from the lichen to the feathermoss, the fen rim, and ending at the fen site. The lichen and the feather moss collars were located along the boardwalk approximately 200 m northeast of the tower. The fen rim and fen site collars were located due east of the lichen and feather moss sites about 150 m and 200 m respectively.

#### For data collected by Dean Moosavi:

- The collars and profile sampling sites near the OBS tower were located as follows: the OBS sampling collars were located along a moisture gradient from feathermoss and lichen sites to the fen rim site and ending with the fen sites. The transect ran approximately east to west and was located about 200 m northeast of the tower at the OBS site.
- The collars and profile sampling sites at the BP tower site were located as follows: the BP sampling sites were located along a low to high moisture gradient that began with the upland lichen site, then the sphagnum moss site, the mire site, and ending at the pond site. The transect ran approximately north to south and was about 60 m from the BP tower.

#### 7.1.2 Spatial Coverage Map

Not available.

#### 7.1.3 Spatial Resolution

Four collars were placed in the ground for each biome type. They were approximately 1 to 2 m apart depending on the site specifics. In Section 7.3.3, the cover type refers to the predominant vegetation in the collar. These are point source measurements at the locations given in Section 7.1.1.

#### 7.1.4 Projection

Not applicable.

#### 7.1.5 Grid Description

Not applicable.

#### 7.2 Temporal Characteristics

#### 7.2.1 Temporal Coverage

Chamber fluxes were measured from 16-May-1994 through 13-Sep-1994. One diurnal study was conducted at eight collars on 08- and 09-Jul-1994 at the BP site.

#### 7.2.2 Temporal Coverage Map

Not available.

#### 7.2.3 Temporal Resolution

The chamber fluxes were measured approximately every 7 days from 23-May-1994 to 20-Sep-1994.

#### 7.3 Data Characteristics

#### 7.3.1 Parameter/Variable

The parameters contained in the data files on the CD-ROM are:

Column Name
SITE_NAME
SUB_SITE
DATE_OBS
TIME_OBS
CO2_FLUX
CH4_FLUX
COVER_TYPE
SOIL_TEMP_1CM
SOIL_TEMP_5CM
SOIL_TEMP_10CM
SOIL_TEMP_20CM
COLLAR_ID
CO2_CONC
CO2_CONC_2
CH4_CONC
CH4_CONC_2
CRTFCN_CODE
REVISION_DATE

#### 7.3.2 Variable Description/Definition

The descriptions of the parameters contained in the data files on the CD-ROM are:

Description		
The identifier assigned to the site by BOREAS, in the format SSS-TTT-CCCCC, where SSS identifies the portion of the study area: NSA, SSA, REG, or TRN; TTT identifies the cover type for the site (999 if unknown); and CCCCC is the		
identifier for site (exactly what it means will vary with site type).		
The identifier assigned to the subsite by BOREAS, in the format GGGGG-IIIII, where GGGGG is the group associated with the subsite instrument (e.g., HYD06 or STAFF), and IIIII is the identifier for the subsite, (often this will		

refer to an instrument). DATE OBS The date on which the data were collected. TIME OBS The Greenwich Mean Time (GMT) when the data were collected. CO2 FLUX Carbon dioxide flux. CH4 FLUX Methane flux. COVER TYPE The dominant species, vegetation, or type of land cover that exists at the location. SOIL TEMP 1CM Soil temperature at 1-cm depth. SOIL TEMP 5CM Soil temperature at 5-cm depth. SOIL TEMP 10CM Soil temperature at 10-cm depth. SOIL TEMP 20CM Soil temperature at 20-cm depth. COLLAR ID A TGB-01 designation for the chamber collar sites, in the form of A-BBB-##, where A = C or M (Crill or Moosavi), BBB denotes the site and possibly microtopographic designation, and ## is a sequention collar number. CO2 CONC CO2 concentration. CO2 CONC 2 CO2 concentration. CH4 CONC CH4 concentration. CH4 CONC 2 CH4 concentration. CRTFCN CODE The BOREAS certification level of the data. Examples are Checked by PI (CPI), Certified by Group (CGR), Preliminary (PRE), and CPI but questionable (CPI-???). The most recent date that the information in the REVISION DATE referenced data base table record was revised.

#### 7.3.3 Unit of Measurement

The measurement units for the parameters contained in the data files on the CD-ROM are:

Column Name	Units
SITE NAME	[none]
SUB SITE	[none]
DATE OBS	[DD-MON-YY]
TIME OBS	[HHMM GMT]
CO2 FLUX	[micromoles][meter-2][second-1]
CH4_FLUX	<pre>[micromoles] [meter-2] [second-1]</pre>
COVER_TYPE	[none]
SOIL_TEMP_1CM	[degrees Celsius]
SOIL_TEMP_5CM	[degrees Celsius]
SOIL_TEMP_10CM	[degrees Celsius]
SOIL_TEMP_20CM	[degrees Celsius]
COLLAR_ID	[none]
CO2_CONC	[parts per million]
CO2_CONC_2	[parts per million]
CH4_CONC	[parts per million]
CH4_CONC_2	[parts per million]
CRTFCN_CODE	[none]
REVISION_DATE	[DD-MON-YY]

#### 7.3.4 Data Source

The sources of the parameter values contained in the data files on the CD-ROM are:

Column Name	Data Source
SITE_NAME	Not applicable
SUB_SITE	Not applicable
DATE_OBS	Investigator
TIME_OBS	Investigator
CO2_FLUX	Calculated by investigator
CH4_FLUX	Calculated by investigator
COVER_TYPE	Determined by investigator
SOIL_TEMP_1CM	Thermocouple
SOIL_TEMP_5CM	Thermocouple
SOIL_TEMP_10CM	Thermocouple
SOIL_TEMP_20CM	Thermocouple
COLLAR_ID	Thermocouple
CO2_CONC	Shimadzu GC-14A
CO2_CONC_2	Shimadzu GC-14A
CH4_CONC	Shimadzu GC-14A
CH4_CONC_2	Shimadzu GC-14A
CRTFCN_CODE	Shimadzu GC-14A
REVISION DATE	Shimadzu GC-14A

## 7.3.5 Data Range

The following table gives information about the parameter values found in the data files on the CD-ROM.

	Minimum	Maximum	Missng			Data
	Data	Data	Data	Data	Detect	Not
Column Name	Value	Value	Value	Value	Limit	Cllctd
SITE_NAME	NSA-BVP-FLXTR	NSA-YJP-FLXTR	None	None	None	None
SUB_SITE	TGB01-FLX01	TGB01-FLX01	None	None	None	None
DATE_OBS	16-MAY-94	13-SEP-94	None	None	None	None
TIME_OBS	0	2316	None	None	None	None
CO2_FLUX	-3.9236111	10.6365741	-999	-888	None	None
CH4_FLUX	-0.0062153	2.4537037	-999	-888	None	None
COVER_TYPE	N/A	N/A	None	None	None	None
SOIL_TEMP_1CM	1	40.3	-999	None	None	Blank
SOIL_TEMP_5CM	2.8	28.4	-999	None	None	Blank
SOIL_TEMP_10CM	-1	23.1	-999	None	None	None
SOIL_TEMP_20CM	-1.9	99.7	-999	None	None	Blank
COLLAR_ID	C-OBS-01	M-OBS-w-16	None	None	None	None
CO2_CONC	363.5	472.5	None	None	None	Blank
CO2_CONC_2	364	504.3	-999	None	None	Blank
CH4_CONC	1.945	2.319	None	None	None	Blank
CH4_CONC_2	2.066	2.262	-999	None	None	Blank
CRTFCN_CODE	PRE	CPI	None	None	None	None
REVISION_DATE	23-AUG-96	14-JAN-97	None	None	None	None

Minimum Data Value -- The minimum value found in the column.

Maximum Data Value -- The maximum value found in the column.

Missng Data Value -- The value that indicates missing data. This is used to indicate that an attempt was made to determine the

parameter value, but the attempt was unsuccessful.

Unrel Data Value -- The value that indicates unreliable data. This is used to indicate that an attempt was made to determine the parameter value, but the analysis personnel deemed the value to be unreliable.

Below Detect Limit -- The value that indicates parameter values below the instrument's detection limits. This is used to indicate that an attempt was made to determine the

indicate that an attempt was made to determine the parameter value, but the analysis personnel determined that the parameter value was below the detection limit of the instrumentation.

Data Not Cllctd -- This value indicates that no attempt was made to determine the parameter value. This usually indicates that BOREAS Information System (BORIS) staff combined several similar but not identical data sets into the same data base table but this particular science team did not measure that parameter.

Blank -- Indicates that blank spaces are used to denote that type of value. N/A -- Indicates that the value is not applicable to the respective column. None -- Indicates that no values of that sort were found in the column.

#### 7.4 Sample Data Record

The following are wrapped versions of data records from a sample data file on the CD-ROM.

```
SITE_NAME, SUB_SITE, DATE_OBS, TIME_OBS, CO2_FLUX, CH4_FLUX, COVER_TYPE, SOIL_TEMP_1CM, SOIL_TEMP_5CM, SOIL_TEMP_10CM, SOIL_TEMP_20CM, COLLAR_ID, CO2_CONC, CO2_CONC_2, CH4_CONC, CH4_CONC_2, CRTFCN_CODE, REVISION_DATE
'NSA-BVP-FLXTR', 'TGB01-FLX01', 02-AUG-94, 1716, -888, -888, 'Mire', 16.4, 15.6, 14.3, 'M-BP-mc-26', 'CPI, '13-JAN-97
```

## 8. Data Organization

#### 8.1 Data Granularity

The smallest unit of data is the CO<sub>2</sub> and CH<sub>4</sub> fluxes measured on a particular date at a particular site.

#### **8.2** Data Format(s)

The Compact Disk-Read-Only Memory (CD-ROM) files contain American Standard Code for Information Interchange (ASCII) numerical and character fields of varying length separated by commas. The character fields are enclosed with single apostrophe marks. There are no spaces between the fields.

Each data file on the CD-ROM has four header lines of Hyper-Text Markup Language (HTML) code at the top. When viewed with a Web browser, this code displays header information (data set title, location, date, acknowledgments, etc.) and a series of HTML links to associated data files and related data sets. Line 5 of each data file is a list of the column names, and line 6 and following lines contain the actual data.

## 9. Data Manipulations

#### 9.1 Formulae

$$\begin{aligned} R_f &= C_{std}/A_{std} \\ C_s &= R_f * A_s \end{aligned}$$

Where:  $R_f$  = Response factor

 $A_{std}$  = Average of 10 standard peak areas  $C_{std}$  = Concentration of the standard  $C_s$  = Concentration of the sample

 $A_s$  = Peak area of sample

CH<sub>4</sub> and CO<sub>2</sub> concentrations were calculated from the average of 10 peak areas of known CH<sub>4</sub> and CO<sub>2</sub> standards. The response factor was calculated as the concentration of the known standard divided by the average of 10 standard peak areas. The peak area of the unknown sample was multiplied by the response factor. Fitting a regression curve to the time series of CH<sub>4</sub> and CO<sub>2</sub> concentrations made the flux calculations. The flux rate of a gas is calculated using the following equation:

 $F = ppmv/min*(P/R*MW)*1/T*V_c/A_c*C$ 

Where:  $F = flux (mg/m^2/d)$ 

P = Chamber pressure (atm) R = 8.2054 x 10-5 m<sup>3</sup> atm/mol/K MW = Molecular weight of the analyte

T = Chamber temperature (K)  $V_c$  = Chamber volume (m<sup>3</sup>)  $A_c$  = Chamber area (m<sup>2</sup>) C = (1000 mg/g\*1440 min/d)

Gases:  $CH_4 = 16 \text{ g/mol}$  $CO_2 = 44 \text{ g/mol}$ 

At sites where oxidation of  $CH_4$  occurs, the flux starts at or near ambient  $CH_4$  concentrations and is depleted below ambient concentrations. If the correlation coefficients were not significantly high, the flux was determined to be below the detection limit and was regarded as a zero flux rate. If there was an efflux of  $CH_4$  was measured at any of these sites, those data were eliminated.

For sites where CH<sub>4</sub> and CO<sub>2</sub> were known to have a positive flux (from the soil into the atmosphere) the regressions of the time series were expected to have an r<sup>2</sup> of greater than 0.85.

If -888 is present in the data set, it indicates that a measurement was taken but discarded for some reason. If -999 is present, then no data were taken.

### 9.1.1 Derivation Techniques and Algorithms

Not applicable.

#### 9.2 Data Processing Sequence

Not applicable.

#### 9.2.1 Processing Steps

The peak areas were taken directly from the HP ChemStation reports from the GC. They were entered into spreadsheets and the concentrations were calculated by the formulas in Section 9.1. The spreadsheets then automatically calculate the flux using the formulas.

#### 9.2.2 Processing Changes

None given.

#### 9.3 Calculations

Not applicable.

#### 9.3.1 Special Corrections/Adjustments

Not applicable.

#### 9.3.2 Calculated Variables

Not applicable.

#### 9.4 Graphs and Plots

None.

#### 10. Errors

#### 10.1 Sources of Error

Field sampling error could account for some error in the concentration of the syringe samples:

- Not flushing the sampling line from the chamber before sampling could cause dilution of the sample with air from the last sampling time.
- Not completely closing the syringes or allowing them to come open during transport will cause dilution from ambient air.
- Placing the chamber down with much force can change the pressure inside the chamber to other than ambient and can affect the mechanisms and processes producing/taking up CH<sub>4</sub> and CO<sub>2</sub>.

Errors such as these would have been written down in the lab/field books, and those data therefore would have been edited out. The analytical precision of the GCs is 0.2% for CH<sub>4</sub> and 1% for CO<sub>2</sub>.

## 10.2 Quality Assessment

#### 10.2.1 Data Validation by Source

None given.

#### 10.2.2 Confidence Level/Accuracy Judgment

None given.

#### 10.2.3 Measurement Error for Parameters

The analytical precision of the GCs is 0.2% for CH<sub>4</sub> and 1% for CO<sub>2</sub>.

#### 10.2.4 Additional Quality Assessments

Not applicable.

#### 10.2.5 Data Verification by Data Center

Data were examined for general consistency and clarity.

#### 11. Notes

#### 11.1 Limitations of the Data

None given.

#### 11.2 Known Problems with the Data

See notes under Section 6.

#### 11.3 Usage Guidance

None given.

#### 11.4 Other Relevant Information

None given.

## 12. Application of the Data Set

The chamber flux data can be used in connection with the tower flux data to determine the  $CH_4$  and  $CO_2$  exchange between the atmosphere and the boreal soils. The soil profile data can be used in comparison with the flux data to determine controls on the fluxes for a certain biome.

#### 13. Future Modifications and Plans

None.

### 14. Software

#### 14.1 Software Description

HP ChemStation.

#### 14.2 Software Access

HP ChemStation can be purchased from Hewlett Packard.

#### 15. Data Access

The TGB-01 NSA CH<sub>4</sub> and CO<sub>2</sub> chamber flux data are available from the Earth Observing System Data and Information System (EOSDIS) Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC).

#### 15.1 Contact Information

For BOREAS data and documentation please contact:

ORNL DAAC User Services Oak Ridge National Laboratory P.O. Box 2008 MS-6407 Oak Ridge, TN 37831-6407 Phone: (423) 241-3952

Fax: (423) 574-4665

E-mail: ornldaac@ornl.gov or ornl@eos.nasa.gov

#### 15.2 Data Center Identification

Earth Observing System Data and Information System (EOSDIS) Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC) for Biogeochemical Dynamics http://www-eosdis.ornl.gov/.

#### 15.3 Procedures for Obtaining Data

Users may obtain data directly through the ORNL DAAC online search and order system [http://www-eosdis.ornl.gov/] and the anonymous FTP site [ftp://www-eosdis.ornl.gov/data/] or by contacting User Services by electronic mail, telephone, fax, letter, or personal visit using the contact information in Section 15.1.

#### 15.4 Data Center Status/Plans

The ORNL DAAC is the primary source for BOREAS field measurement, image, GIS, and hardcopy data products. The BOREAS CD-ROM and data referenced or listed in inventories on the CD-ROM are available from the ORNL DAAC.

## 16. Output Products and Availability

## 16.1 Tape Products

None.

#### 16.2 Film Products

None.

#### 16.3 Other Products

These data are available on the BOREAS CD-ROM series.

### 17. References

### 17.1 Platform/Sensor/Instrument/Data Processing Documentation

Instruction Manual: GC-14A. Shimadzu Corporation, Kyoto, Japan.

#### 17.2 Journal Articles and Study Reports

Newcomer, J., D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers, eds. 2000. Collected Data of The Boreal Ecosystem-Atmosphere Study. NASA. CD-ROM.

Sellers, P. and F. Hall. 1994. Boreal Ecosystem-Atmosphere Study: Experiment Plan. Version 1994-3.0, NASA BOREAS Report (EXPLAN 94).

Sellers, P. and F. Hall. 1996. Boreal Ecosystem-Atmosphere Study: Experiment Plan. Version 1996-2.0, NASA BOREAS Report (EXPLAN 96).

Sellers, P., F. Hall, and K.F. Huemmrich. 1996. Boreal Ecosystem-Atmosphere Study: 1994 Operations. NASA BOREAS Report (OPS DOC 94).

Sellers, P., F. Hall, and K.F. Huemmrich. 1997. Boreal Ecosystem-Atmosphere Study: 1996 Operations. NASA BOREAS Report (OPS DOC 96).

Sellers, P., F. Hall, H. Margolis, B. Kelly, D. Baldocchi, G. den Hartog, J. Cihlar, M.G. Ryan, B. Goodison, P. Crill, K.J. Ranson, D. Lettenmaier, and D.E. Wickland. 1995. The boreal ecosystem-atmosphere study (BOREAS): an overview and early results from the 1994 field year. Bulletin of the American Meteorological Society. 76(9):1549-1577.

Sellers, P.J., F.G. Hall, R.D. Kelly, A. Black, D. Baldocchi, J. Berry, M. Ryan, K.J. Ranson, P.M. Crill, D.P. Lettenmaier, H. Margolis, J. Cihlar, J. Newcomer, D. Fitzjarrald, P.G. Jarvis, S.T. Gower, D. Halliwell, D. Williams, B. Goodison, D.E. Wickland, and F.E. Guertin. 1997. BOREAS in 1997: Experiment Overview, Scientific Results and Future Directions. Journal of Geophysical Research 102(D24): 28,731-28,770.

## 17.3 Archive/DBMS Usage Documentation None.

## 18. Glossary of Terms

None given.

## 19. List of Acronyms

AES - Atmospheric Environment Services

ASCII - American Standard Codes for Information

BOREAS - BOReal Ecosystem-Atmosphere Study

BORIS - BOREAS Information System
BP - Beaver Pond site, NSA

CD-ROM - Compact Disk-Read-Only Memory

CGR - Certified by Group

CMDL - Climate Monitoring and Diagnostics Laboratory

CPI - Certified by PI

CPI-??? - Certified but questionable

DAAC - Distributed Active Archive Center

ECD - Electron Capture Detector
EOS - Earth Observing System

EOSDIS - EOS Data and Information System

FID - Flame Ionization Detector

GC - Gas Chromatograph

GIS - Geographic Information System

GMT - Greenwich Mean Time

GSFC - Goddard Space Flight Center

HP - Hewlett Packard

HTML - HyperText Markup Language NAD83 - North American Datum of 1983

NASA - National Aeronautics and Space Administration NOAA - National Oceanic and Atmospheric Administration

NSA - Northern Study Area
OBS - Old Black Spruce
OJP - Old Jack Pine

ORNL - Oak Ridge National Laboratory PANP - Prince Albert National Park

PRE - Preliminary

SSA - Southern Study Area

TCD - Thermal Conductivity Detector
TGB - Trace Gas Biogeochemistry
URL - Uniform Resource Locator
YJP - Young Jack Pine site, NSA

#### 20. Document Information

#### **20.1 Document Revision Dates**

Written:

Last Updated: 08-Jun-1999

#### 20.2 Document Review Dates

BORIS Review: 21-Oct-1997

Science Review:

#### 20.3 Document ID

#### 20.4 Citation

When using these data, please contact the investigators listed in Section 2.3 and cite any relevant papers in Section 17.2.

If using data from the BOREAS CD-ROM series, also reference the data as:

Crill, P.M., "Magnitude and Control of Trace Gas Exchange in Boreal Ecosystems." In Collected Data of The Boreal Ecosystem-Atmosphere Study. Eds. J. Newcomer, D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers. CD-ROM. NASA, 2000.

#### Also, cite the BOREAS CD-ROM set as:

Newcomer, J., D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers, eds. Collected Data of The Boreal Ecosystem-Atmosphere Study. NASA. CD-ROM. NASA, 2000.

#### 20.5 Document Curator

#### 20.6 Document URL

## REPORT DOCUMENTATION PAGE

Form Approved

OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

Barto riigittay, baito 1201, 711111gton, 177 22202 10	, <b>-</b>	Budget, Tupertrent Headerton	r reject (e.e., cree), tradimigton, Be 200001
1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE 3. REPORT TYPE AND DATES COVERED			
	November 2000	Techn	ical Memorandum
<b>4. TITLE AND SUBTITLE</b> Technical Report Series on the Boreal Ecosystem-Atmosphere Study (BOREAS) BOREAS TGB-1 NSA ${\rm CH_4}$ and ${\rm CO_2}$ Chamber Flux Data			923
6. AUTHOR(S)			RTOP: 923-462-33-01
Patrick Crill and Ruth K. Var	ner		
Forrest G. Hall and Sara K. C	Conrad, Editors		
7. PERFORMING ORGANIZATION NAM	E(S) AND ADDRESS (ES)		8. PEFORMING ORGANIZATION REPORT NUMBER
Goddard Space Flight Center			
Greenbelt, Maryland 20771			2000-03136-0
9. SPONSORING / MONITORING AGE	. ,	(ES)	10. SPONSORING / MONITORING AGENCY REPORT NUMBER TM—2000–209891
National Aeronautics and Space Administration Washington, DC 20546-0001			Vol. 219
Trainington, DO 200 TO 0001			VOI. 217
11. SUPPLEMENTARY NOTES			
P. Crill and R.K. Varner: Uni	versity of New Hampshire	e, Durham;	
S.K. Conrad: Raytheon ITSS			
12a. DISTRIBUTION / AVAILABILITY STA	ATEMENT		12b. DISTRIBUTION CODE
Unclassified-Unlimited			
Subject Category: 43			
Report available from the NASA	A Center for AeroSpace Info	ormation,	
7121 Standard Drive, Hanover,	MD 21076-1320. (301) 62	1-0390.	
13. ABSTRACT (Maximum 200 words)			
EL DODE LO EGO 1	1 1 (CIT) 1	1 1: :1 :6	O \ 1 1 1 1 0

The BOREAS TGB-1 team made methane ( $\mathrm{CH_4}$ ) and carbon dioxide ( $\mathrm{CO_2}$ ) dark chamber flux measurements at the NSA-OJP, NSA-OBS, NSA-BP, and NSA-YJP sites from 16-May-1994 through 13-Sep-1994. Gas samples were extracted approximately every 7 days from dark chambers and analyzed at the NSA lab facility. The data are provided in tabular ASCII files.

14. SUBJECT TERMS BOREAS, trace gas biog	15. NUMBER OF PAGES 16 16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT $UL$